

Figure 1-1. Road Map to Exposure Factor Recommendations

EXPOSURE ROUTE	EXPOSURE FACTOR	POPULATION	VOLUME	CHAPTER	RECOMMENDATIONS/ RATINGS TABLE PAGE NOS.
Ingestion	Drinking Water Intake Rate	Adults Children Pregnant Women High Activity	ı	3.	3-25/3-31
	Fruit and Vegetable Intake Rate	Various Demographic Groups — Age,	II	9.	9-8/9-40,9-41
	- Meat and Dairy Intake Rate	Region, Season, Urbanization, Race	II	11.	11-6/11-26,11-27
	Homegrown Foods		II	12.	12-9/12-77
	Breast milk Intake Rate	Nursing Infants	II	13.	13-8/13-14
	Fish and Shellfish Intake Rate	General Population	II	10.	10-49/10-57
		Freshwater Recreational	II	10.	10-51/10-59
		Marine Recreational	II.	10.	10-50/10-58
	Soil Intake Rate	Subsistence Typical Children Adults Pica Children	I	10. 4.	10-51/10-60 4-19/4-23
Inhalation ————————————————————————————————————	- Inhalation Rate	Adults Children High Activity	ı	5.	5-21/5-21
Dermal —	- Skin Surface Area		I	6.	6-8/6-23
	- Soil Adherence	— General Population	1	6.	6-8/6-24
(All Routes) Human Characteristics	— Body Weight	Adults Children	1	7.	7-7/7-9
	Lifetime —		1	8.	8-1/8-3
(All Routes) Activity Factors	- Activity Patterns	Adults Children	Ш	14.	1-69/1-72
	— Occupational Mobility —		III	14.	1-71/1-75
	Population Mobility —		III	14.	1-71/1-44
(All Routes) Consumer Product Use	- Frequency of Use	Adults	III	45	15-5
	- Amount Used -	Adults	III	15.	15-5
(All Routes) Reference House	- Water Use - Air Exchange Rates - House Volumes - Building Characteristics	General Population	III	16.	16-25/16-32,16-33









Figure 6-1. SA/BW Distributions for Infants, Adults, and All Ages Combined Source: Phillips et al., 1993.







Figure 6-2. Frequency Distributions for the Surface Area of Men and Women Source: Murray and Burmaster, 1992.

Area in m², n=5,000, LHS



* Participants could list more than one month.

Figure 10-1. Seasonal Fish Consumption: Wisconsin Chippewa, 1990




Figure 10-2. Peak Fish Consumption: Wisconsin Chippewa, 1990

Source: Peterson et al., 1994.




Figure 14-1. Distribution of Individuals Moving by Type of Move: 1991-92

Source: U.S. Bureau of the Census, 1993






Figure 16-1 Elements of Residential Exposure



COMMON RETURN LAYOUT



BALANCED SUPPLY and RETURN LAYOUT








Figure 16-3. Configuration for Residential Forced-air Systems











BEDROOM VERSUS REMAINDER OF HOUSE



Bedroom Volume = 35 m³ Remainder of House Volume = 373 m³

LIVING ROOM VERSUS REMAINDER OF HOUSE



Living Room Volume = 60 m³ Remainder of House Volume = 348 m³

Figure 16-7. Characteristic Volumes and Air Flow Rates for Two-Zone Situations